


# Targeted Temperature Management in Trauma



1. see ACLS protocols in Circulation 2005; 112(24):IV84-IV88.
2. Although many institutions routinely cool patients with lung injury, shock, and moderate coagulopathy, data is lacking. It is not known whether hypothermia has neuroprotective effects if therapy is delayed beyond 6 hours after ROSC.
3. TH is probably safe in combination with PTCA/PCI, intra-aortic balloon pump placement, and thrombolytics. See Crit Care Med 2006;34(6):1865-73 and Acta Anaesthesiol Scand 2007;51:137-42.
4. Sirvent JM et al. Protective Effect of Intravenously Administered Cefuroxime Against Nosocomial Pneumonia in Patients with Structural Coma. Am J Respir Crit Care Med 1997; 155: 1729-34.
5. Crompton EM et al. Meta-Analysis of Therapeutic Hypothermia for Traumatic Brain Injury in Adult and Pediatric Patients. Critical Care Medicine. 2017; 45(4); 575-579.
6. Hsu CH et al. EAST Multicenter Trial on Targeted Temperature Management for Hanging Induced Cardiac Arrest. J Trauma Acute Surg. 2018; 8(1); 37- 47.
7. Yang Z, Song Z, Hou M. Target Temperature Management Versus Normal Temperature Management for Cardiac Arrest After Traumatic Brain Injury Patient: A Meta-Analysis and Systemic Review. Ther Hypothermia Temp Manag. 2022 Sep;12(3):139-145.. Epub 2022 Aug 1. PMID: 35914088.